

Impact of Watershed Management on Irrigated Areas

S.N.	Source of irrigation	Irrigated Area (Ha)		
		Before	After Treatment	Change
1	Earthen Embankment	19.50	977.60	958.10
2	Pond	11.60	451.20	439.60
3	Natural <i>Nala</i>	34.90	75.20	40.30
	Total	66.00	1,504.00	1,438.00



Crop development at Majhgawan — Wheat.

Impact of Watershed Management on Cropped Area

	Sl. No.	Crop	Cropped Area (ha)		
			Before	After treatment	Change
	1	Cereals	546.26	825.5	279.24
	1.1	Paddy	170.40	247.65	77.25
	1.2	Sorgham	108.94	99.06	-9.88
	1.3	Wheat	178.50	396.24	217.74
	1.4	Barley	88.42	82.55	-5.87
	2	Pulses	242.40	450.2	207.8
	2.1	Pigeon Pe	111.80	90.04	-21.76
	2.2	Gram	131.60	360.16	228.56
	3	Oilseeds	169.45	228.3	58.85
	3.1	Mustard	84.85	159.81	74.96
	3.2	Linseed	51.20	45.66	-5.54
	3.3	Til	33.40	22.83	-10.57
	Total		958.11	1504	545.89